

Instability in the Linkage of Topological Spaces Due to Background Ghosts

Deep Bhattacharjee

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

April 12, 2023

Instability in the linkage of topological spaces due to background ghosts

Deep Bhattacharjee[^]

The linkage of geometric structures in certain dimensions with each other provided the 'link node' is stable can in essence establish the Hopf fibrations in the homotopy theory but under certain 'oscillating backgrounds' or 'topological ghosts' as termed in this paper might destroy the link and make each 'geometric structure' independent on their own.

```
Ghosts – Factor – Nodes
```

Taking a background space *B* having the boundary ∂ for a coherent norm of boundary space ∂B there can be a relation through nodes *N* acting on a stable parameter γ on the before mentioned background space ∂B which in essence is a complex topological space denoted as T^* for a mapping parameter \star such that the nodes or links *N* can take two values for $\partial B^{[1-4]}$,

$\begin{cases} stable parameter \gamma \\ unstable parameter \gamma_{\times} \end{cases}$

Denoting # as an affine parameter for the periodicity ρ determining the oscillation factor ε to give the notions^[5-6],

$$\# = \sum_{\rho=1}^{<\infty} \varepsilon_{\rho} \Longrightarrow \star : \begin{cases} \gamma \stackrel{\hookrightarrow}{\to} \gamma_{\times} \\ \gamma_{\times} \stackrel{\hookrightarrow}{\to} \gamma \end{cases} \exists \begin{cases} \star_{0} : \gamma \longrightarrow \gamma_{\times} \\ \star_{1} : \gamma_{\times} \longrightarrow \gamma \end{cases}$$

Over a dependency parameter ∇ with a sub ∇_f such that f is the frequency parameter which takes two values^[7-10],

 $f \ni \begin{cases} f_{\Pi 1} \exists 1 \text{ is the increase in order of frequencies via multiplicity} \\ f_{\Pi 0} \exists 0 \text{ is the decrease in order of frequencies via no multiplicity} \\ \begin{cases} \forall \text{stability and instability conditions} \\ \downarrow \\ N \ni \begin{cases} N_{++} := \partial B_{\star_0} \xrightarrow{f_{\Pi 1}} \partial B_{\star_1} \\ N_{\times \times} := \partial B_{\star_1} \xrightarrow{f_{\Pi 0}} \partial B_{\star_1} \end{cases} \end{cases}$

[^]INS Research, Department of Geometry and Topology, India

[^]Electro – Gravitational Space Propulsion Laboratory, India-

[^]CXAI Technologies Ltd., Cyprus+

^{^+}RA on Computations of Theoretical Actual Intelligence (AcI) Formalism

[^]Former affiliations with AATWRI Aerospace & Defence Research Directorate, India

[^]The author of this paper holds no conflict of interest as related to this paper [^]Author electronic address: itsdeep@live.com

Thus, considering two topological structures T_1 and T_2 which can be extended up to T_n for a finite order of $n < \infty$; this can be represented via the dependency parameter ∇_f this can be represented via^[11,12],

$$\eta_{\nu}{}^{\zeta_{S^i}}: \quad T_1 \xrightarrow{\nabla_f} T_2 \ \dots \xrightarrow{\nabla_f} T_n$$

Hopf connectivity can not be possible if in ζ_{S^i} for ζ as the mapping with S^i representing the sphere with the suspension η_{ν} acting on ζ_{S^i} for $\nu \equiv +$ in case of suspension with no connectivity or fibrations with the background oscillations and $\nu \equiv -$ the structural suspensions without background oscillations nulifying the linkage of spheres S^i for the various dimensions represented by *i* if and only if ∇_f represents with the fibration parameter $\mathcal{F}^{[\text{Fig. 13, 14-18}]}$,

$$\mathcal{F}_{\epsilon} \begin{cases} \mathcal{F}_{\pm} \Longrightarrow no \ fibrations \\ else \\ \mathcal{F}_{c} \Longrightarrow \ fibrations \end{cases} \exists \{\pm, c\} \subset \epsilon \xrightarrow{\eta_{\nu}} \end{cases}$$



[#]Null set ϕ is the element of every set.

The ghosts which here termed as instabilities arise out of the background oscillations of the topological space T^* where the instabilities arise out of two factor and affects the third which is the boundary of the geometric structure ∂B which links to each other in a way of dimensions that can be same or cannot such that this ∂B can be formulated via the structure dependent on the geometric spaces having the form^[19-21],

	inequalities
$(\partial B)_d$	$d, \bar{d} = \bar{\bar{d}} \{ S^0 \hookrightarrow S^1 \hookrightarrow S^1$
$(\partial B)_{\bar{d}}$	$d \neq \bar{d} \neq \bar{\bar{d}} \notin \bar{\bar{d}} \begin{cases} S^1 \hookrightarrow S^3 \hookrightarrow S^2 \\ S^3 \hookrightarrow S^7 \hookrightarrow S^4 \\ S^7 \hookrightarrow S^{15} \hookrightarrow S^8 \end{cases}$
$(\partial B)_{\bar{d}}$	

The two factor for which the instability arises which has been termed as ghost on the background space $T^* \exists (\partial B)_{d,\bar{d},\bar{d},\bar{d}}$ exists on T^* are^[22,23],

- *N* the node of the linage between d, \bar{d}, \bar{d} acting on *S*.
- ρ is the oscillation of *N* implying *N* is a function of ρ as $N(\rho) = \eta_{\nu}$ where ν_{ν} vanishes for Hopf fibration where frequency *f* won't act on ρ ,

$$\begin{array}{ccc} & \exists f & & \overbrace{ \ }^{\{\phi\},\mathcal{F}_{c}} & \rho \mid \\ & & \nu \text{ vanishes in } \eta_{\nu} \zeta_{s^{i}} \\ \bullet & else \ f & \xrightarrow{\rightarrow} \rho \ \text{ for } & \underbrace{ \{N_{++} \ N_{\times \times} \},\mathcal{F}_{\pm}}_{\nu=+,-in \ \eta_{\nu}} \zeta_{s^{i}} \implies \text{ no fibration} \end{array}$$

REFERENCES:

- **1.** Zeuthen, H. G., Kleiman, S. L., & Thorup, A. (1991). *Enumerative Algebraic Geometry: Proceedings of the 1989 Zeuthen Symposium*. American Mathematical Soc.
- **2.** Lebarz, & Hervier. (2012). *Enumerative Geometry and Classical Algebraic Geometry*. Springer Science & Business Media.
- **3.** Bhattacharjee, D. (2022b). Suspension of structures in two-dimensional topologies with or without the presence of $g \ge 1$ genus deformations for canonical $2^2\eta$ stabilizer points. *Research Square*. https://doi.org/10.21203/rs.3.rs-1798323/v1
- **4.** Bhattacharjee, D., Samal, P., Bose, P. N., Behera, A. K., & Das, S. (2023, April 5). Suspension η for β bundles in ± 1 geodesics in $g \ge 1$ genus creations for loops for a Topological String Theory Formalism. https://doi.org/10.36227/techrxiv.22339732.v1
- 5. Mendelson, B. (1990). Introduction to Topology. Courier Corporation.
- Bhattacharjee, D. (2022d). Non-commutativity Over Canonical Suspension η for Genus g ≥1 in Hypercomplex Structures for Potential pφ. Asian Research Journal of Mathematics, 332–341. https://doi.org/10.9734/arjom/2022/v18i11607

- **7.** Arkowitz, M. (2011). *Introduction to Homotopy Theory*. Springer Science & Business Media.
- Bhattacharjee, D (2022): An outlined tour of geometry and topology as perceived through physics and mathematics emphasizing geometrization, elliptization, uniformization, and projectivization for Thruston's 8-geometries covering Riemann over Teichmuller spaces. TechRxiv. Preprint. https://doi.org/10.36227/techrxiv.20134382.v1
- 9. Adams, J. F., & Atiyah, M. (1966). *K*-THEORY AND THE HOPF INVARIANT. *Quarterly Journal of Mathematics*, *17*(1), 31–38. https://doi.org/10.1093/qmath/17.1.31
- Bhattacharjee, Deep; Sadhu, Riddhima; Behera, Ashis Kumar; Roy, Sanjeevan Singha; Das, Saptashaw (2023): KK Theory and K Theory for Type II Strings Formalism. TechRxiv. Preprint. https://doi.org/10.36227/techrxiv.21805215.v1
- **11.** Adams, J. F. (1960). On the Non-Existence of Elements of Hopf Invariant One. *Annals of Mathematics*, 72(1), 20. https://doi.org/10.2307/1970147
- Bhattacharjee, D. (2022h). Establishing Equivariant Class [O] for Hyperbolic Groups. Asian Research Journal of Mathematics, 362–369. https://doi.org/10.9734/arjom/2022/v18i11615
- **13.** Clark, N. S. (2017, February 22). *MATHEMATICS NATURE*. Pinterest. https://www.pinterest.com/pin/208713763962061332/
- 14. Bhattacharjee, D. Establishing equivalence among hypercomplex structures via Kodaira embedding theorem for non-singular quintic 3-fold having positively closed (1,1)-form Kähler potential i2⁻¹∂∂*p, 10 May 2022, PREPRINT (Version 1) available at Research Square https://doi.org/10.21203/rs.3.rs-1635957/v1
- 15. Bhattacharjee, D., Generalization of Quartic and Quintic Calabi Yau Manifolds Fibered by Polarized K3 Surfaces, 16 August 2022, PREPRINT (Version 1) available at Research Square https://doi.org/10.21203/rs.3.rs-1965255/v1
- 16. Bhattacharjee, D., Singha Roy, S., Behera, A. K. Relating Enrique surface with K3 and Kummer through involutions and double covers over finite automorphisms on Topological Euler–Poincaré characteristics over complex K3 with Kähler equivalence, 31 August 2022, PREPRINT (Version 1) available at Research Square https://doi.org/10.21203/rs.3.rs-2011341/v1
- **17.** Kochman, S. O. (2006). *Stable Homotopy Groups of Spheres: A Computer-Assisted Approach*. Springer.
- **18.** Ravenel, D. C. (2003). *Complex Cobordism and Stable Homotopy Groups of Spheres*. American Mathematical Soc.
- **19.** Bhattacharjee, D. (2022c). Mathematical Inequalities over Hypercomplex Structures and Infinities with Related Paradoxes. *EasyChair Preprint No. 8377*. https://easychair.org/publications/preprint/Nh9w
- **20.** Switzer, R. M. (2017). *Algebraic Topology Homotopy and Homology*. Springer.
- **21.** Adams, J. F., & Adams, J. F. (1974). *Stable Homotopy and Generalised Homology*. University of Chicago Press.
- 22. Bhattacharjee, N. D., Roy, N. R., & Sadhu, N. J. (2022b). HOMOTOPY GROUP OF SPHERES, HOPF FIBRATIONS AND VIL-LARCEAU CIRCLES. *EPRA International Journal* of Research & Development, 57–64. https://doi.org/10.36713/epra11212
- 23. May, J. P. (1992). Simplicial Objects in Algebraic Topology. University of Chicago Press.